



Supplement of

The effect of marine aggregate parameterisations on nutrients and oxygen minimum zones in a global biogeochemical model

Daniela Niemeyer et al.

Correspondence to: Daniela Niemeyer (dniemeyer@geomar.de)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

This file provides additional figures to the manuscript ‘The effect of marine aggregate parameterisations on nutrients and oxygen minimum zones in a global biogeochemical model’. It contains the following figures:

Figure S1: Global maps of b for noAgg^{ECCO1.0} (a) (same as Fig. 1a) and simulation BUR (b) from Kriest and Oschlies (2013), where remineralisation does not depend on oxygen.

Figure S2: Global map of b for the four best model simulations with regard to the sum of J_{RMSE} and J_{OMZ} in ECCO1.0: (a) simulation #14; (b) simulation #17; (c) simulation #28; (d) simulation #29).

Figure S3: As Fig. 2, but for $O_2 \leq 30 \text{ mmol m}^{-3}$.

Figure S4: Profiles of average nutrient and oxygen concentrations for the eastern tropical Pacific (upper panels) and globally (lower panels).

Figure S5: Simulated particle abundance (in $\# \text{ L}^{-1}$) in the surface layer (upper), for the 23°W section (lower left) and 151°W section (lower right) as a depth-profile for a particle size range of 0.14 to 16.88 mm diameter, following Kiko et al. (2017).

Supplement Figures




Figure S1: Global maps of b for noAgg^{ECCO1.0} (a) (same as Fig. 1a) and simulation BUR (b) from Kriest and Oschlies (2013), where remineralisation does not depend on oxygen.




Figure S2: Global map of b for the four best model simulations with regard to the sum of J_{RMSE} and J_{OMZ} in ECCO1.0: (a) simulation #14; (b) simulation #17; (c) simulation #28; (d) simulation #29.




Figure S3: As Fig. 2, but for $\text{O}_2 \leq 30 \text{ mmol m}^{-3}$.




Figure S4: Profiles of average nutrient and oxygen concentrations for the eastern tropical Pacific (upper panels) and globally (lower panels).




Figure S5: Simulated particle abundance (in # L^{-1}) in the surface layer (upper) and for the 23°W section (lower left) and 151°W section (lower right) as a depth-profile for a particle size range of 0.14 to 16.88 mm diameter, following Kiko et al. (2017).