



The SUMup dataset: Compiled measurements of surface mass balance components over ice sheets and sea ice with preliminary analysis over Greenland

Lynn Montgomery¹, Lora Koenig², Patrick Alexander^{3,4}

5 ¹Department of Atmospheric and Oceanic Science, University of Colorado, Boulder, CO, USA

²National Snow and Ice Data Center, University of Colorado, Boulder, CO, USA

³Lamont Doherty Earth Observatory, Columbia University, Palisades, NY, USA

⁴NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY 10025, USA

10 *Correspondence to:* lynn.montgomery@colorado.edu

Abstract

Increasing atmospheric temperatures over ice cover affects surface processes including melt, snowfall and snow density. Here, we present the SUMup dataset, a standardized dataset of Arctic and Antarctic observations of surface mass balance components. The July 2017 SUMup dataset consists of three subdatasets, snow/firn density (doi:10.18739/A26D6F), snow accumulation on land ice (doi:10.18739/A2XX0V), and snow depth on sea ice (doi:10.18739/A22Q35), to monitor change and improve estimates of surface mass balance. The measurements in this dataset were compiled from field notes, papers, technical reports, and digital files. SUMup is a compiled, community-based dataset that can be and has been used to evaluate modeling efforts and remote sensing retrievals. Measurements in the dataset are sporadic in time and have spatial gaps, however, they likely constitute the largest set of field measurements compiled, standardized and publicly available. Analysis of the dataset shows that Greenland ice sheet density measurements in the top 1 m do not show a strong relationship with annual temperature. At Summit Station, Greenland, accumulation and surface density measurements vary seasonally with lower values during summer months. The SUMup dataset is a dynamic, living dataset that will be updated and expanded for community use as new measurements are taken and new processes are discovered and quantified.

Copyright Statement

We agree to the Creative Commons Attribution 4.0 License and the Copernicus Publications rights.

1. Introduction and Background

30 Earth's polar regions are warming at an accelerated rate. As increased air temperatures, and associated feedbacks with radiative heating, persist the ice cover is changing particularly at the ice/atmosphere interface (e.g. Vaughan and others, 2003; Serreze and Francis, 2006; Hall and others, 2013). This change is evident by declining sea ice extent (e.g. Ritcher-Menge and others, 2016) and the recent acceleration of mass loss from the Greenland Ice Sheet (GrIS) and Antarctic ice sheets (AIS), (e.g. Velicogna and others, 2014; Shepherd and others, 2012), that contributed ~11 mm to global sea levels between 1992 and 2011 (Shepherd and others, 2012). Surface change is particularly evident over the GrIS where surface mass balance processes, dominated by melt, now account for more than half of mass loss through surface melting and decreasing surface albedo (e.g. van Angelen and others, 2013; van den Broeke and others, 2009; Enderlin and others, 2014). To understand this change, and the processes driving the change, it is vital to compile measurements from the past and present, and continue to collect them in the future.

40

In 2012, at the Surface Mass Balance and Snow on Sea Ice Working Group (SUMup) meeting, the modeling and remote sensing communities clearly stated to observationalists that the lack of easy to access, standardized, in-situ measurements hindered scientific achievement. A public, annual, decadal, standardized time-series of measurements was recommended (Koenig and others, 2013). Modelling and remote sensing studies require validation measurements (e.g. Fettweis and others,



45 2017; Arthern and others, 2006; Burgess and others, 2010; Kuipers Munneke and others 2015; Koenig and others, 2016), ideally at their same spatial and temporal resolutions and over large sections of ice sheets or sea ice, which are difficult for an individual researcher to compile. Today, most field observations for validation are dispersed across multiple data centers/datasets in differing formats. While some previous Arctic and Antarctic studies have compiled large sets of measurements, generally accumulation measurements (e.g. Mock 1967a; Mock 1967b; Ohmura and Reeh, 1999; Vaughan and
50 others, 1999; Arthern and others, 2006), not all of this data are public, formatted consistently, or has the necessary temporal or spatial resolution for recent models.

Here, we present the July 2017 SUMup dataset and its three subdatasets: density, accumulation, and snow depth on sea ice. This data paper serves to fully describe the dataset and includes preliminary analysis of the data over the GrIS demonstrating
55 how this dataset increases our knowledge of surface mass balance processes by compiling previously dispersed measurements into a standardized dataset. Uses of SUMup include model validation, remote sensing validation and algorithm development, and long-term monitoring efforts.

2. The SUMup Dataset

2.1 Overview

60 The SUMup dataset is an expandable, community-based dataset of field measurements of surface mass balance components that is consistent in format, properly described through metadata, and publically available. The July 2017 SUMup dataset contains three subdatasets that consist of measurements of snow/firn density (doi:10.18739/A26D6F), snow accumulation on land ice (doi:10.18739/A2XX0V), and snow depth on sea ice (doi:10.18739/A22Q35). The SUMup dataset is a living document, meant to be expanded as new measurements are taken or previous measurements discovered. The current release
65 of the July 2017 SUMup dataset expands the previous two smaller releases of July 2013 and July 2015 (<https://neptune.gsfc.nasa.gov/csb/index.php?section=267>). The measurements compiled in SUMup are from the polar regions and most date from 1950 to present day. (~2% of the accumulation subdataset predates 1950 and these measurements are included to keep complete records from ice cores.)

70 Figure 1 shows the locations of measurements represented by the July 2017 SUMup dataset excluding ~40 snow depth on sea ice locations off the coast of Finland in the Baltic sea. Density and accumulation measurements are often co-located over the ice sheets where ice cores were collected (Fig. 1).

Figure 1

75 2.2 Sources

SUMup data were collected, formatted and compiled primarily through two methods: 1) searching data archives that traditionally host cryospheric data which included Pangaea (<https://www.pangaea.de/>), the Arctic Data Center (<https://arcticdata.io/>), NOAA's National Climate Data Center (<https://www.ncdc.noaa.gov/>) and the National Snow and Ice Data Center (<https://nsidc.org/>) and 2) by asking members of the cryospheric community to contribute field measurements.
80 Keyword searches for the first method includes searching for the words “density”, “accumulation”, and “snow depth on sea ice”. We look specifically for annually resolved accumulation measurements if possible. Various data types are compiled into the SUMup dataset including hand-written notes, technical reports, and digital files. Each measurement in the SUMup dataset contains a citation to the original source of the data. Based on keyword searches, data for this release, July 2017, should include most relevant data available in the data archival centers listed above posted before April 2016 with a focus on
85 Greenland.

New and unique data sources are included in the SUMup dataset. Notably, the snow density subdataset includes snow pit data from Carl Benson's Greenland traverses in the early 1950's that, to our knowledge, have never before been digitized into a



90 dataset, and include data from 1955 that previously had not been digitally scanned (Benson, 2013; Benson, 2017). The 1955
notebooks are only archived in the National Snow and Ice Data Center paper archives. The SUMup dataset also includes snow
accumulation measurements from Summit Station, Greenland's stake network called the Bamboo Forest (Dibb and others,
2004) and corresponding density measurements at monthly temporal resolution (Dibb and others, 2007). Additionally, more
widely used data sources are included, such as, US International Trans-Antarctic Scientific Expedition (US ITASE, Mayewski
95 et al, 2013) ice cores, the Program for Arctic Regional Climate Assessment (PARCA, Mosley-Thompson et al, 2001) ice cores
and The Greenland Inland Traverse (GrIT, Hawley et al, 2014) snow pits and ice cores. Section 2.4 provides more details on
the specific sources for each of the three subdatasets including the complete list of all citations.

2.3 Contributing to the Dataset

The SUMup dataset will continue to expand on an annual basis as new measurements are taken and/or old measurements are
discovered. Beyond expanding the current subdatasets, we expect to add additional subdatasets on surface mass balance
100 processes which may include, but are not limited to, snow/ice albedo, snow temperature, and short-wave/long-wave radiation
measurements. The community is encouraged to contribute data or suggest missing data sources/types to add to SUMup by
contacting the authors directly.

2.4 Structure and MetaData

Each measurement contains common variables including the date taken, latitude, longitude, surface elevation if on land, the
105 measurement itself, error associated with the measurement, the method in which the measurement was taken, and a citation
for which the measurement can be sourced back to. By convention, negative latitudes represent south and negative longitudes
represent west. For data that did not specify a specific date, but provided only the year, the date was entered as 'yyyy0000'.
A consistent fill value of -9999 was used for unknown or unmeasured parameters.

110 If any of the original data/metadata were unclear or nonexistent, the original author of the data was contacted to clarify
inconsistencies or questions so the SUMup metadata is complete. In total, only ten density measurements from Mayewski and
Whitlow (2016) were not included in the final SUMup dataset. These measurements were either recorded as 0 g/cm^3 or had
density values of over 1 g/cm^3 which would both be impossible for clean ice. Specific details on measurement methods and
citations for each subdataset are included in the SUMup metadata files hosted at the Arctic Data Center and described below.

115 2.4.1 Snow Density

The snow/firn density subdataset of SUMup is the largest, containing over 830,000 point measurements of density at different
depths (Fig. 1). Table 1 describes the parameters for each density measurement. The measurement methods include density
cutters of different sizes (generally from $100 - 1000 \text{ cm}^3$) used in snow pits, gravitational methods used on ice core sections,
neutron-density methods performed in boreholes, X-ray microfocus computer tomography performed on snow samples,
120 gamma-ray attenuation in boreholes and pycnometers used on snow samples. The majority of the observations (~94%) come
from Greenland ice cores or snow pits (Ohmura, 1991; Ohmura, 1992; Alley, 1999; Bolzan and Strobel, 1999(a-g); Miller and
Schwager, 2000(a,b); Wilhelms, 2000(a,b,c,d); Bolzan and Strobel, 2001(a,b); Mosley-Thompson and others, 2001; Bales and
others, 2001; Conway, 2003; Dibb and Fahnestock, 2004; Dibb and others, 2007; Baker and others, 2009; Chellman and others,
2009; Benson, 2013; Miège and others, 2013; Hawley and others, 2014; Koenig and others, 2014; Mayewski and Whitlow,
125 2016; Schaller and others, 2016; Benson, 2017). Antarctic data comprises ~6% of the snow density subdataset and is
predominantly from ice cores (Mayewski et al, 2013; Lewis et al, 2011; Brucker and Koenig, 2011; Medley et al, 2013; Albert,
2007; Kreutz et al, 2011). The depth of the density measurements were recorded using two different methods, either the top
and bottom depth or a midpoint depth. While a midpoint can be determined uniquely from the top and bottom depths the top
and bottom cannot always be determined from the midpoint and researchers need to determine how to standardize or interpolate
130 the depths for their specific applications.

Table 1



2.3.2 Snow Accumulation on Land Ice

135 The snow accumulation on land ice subdataset of SUMup contains over 230,000 data points (Fig. 1). Table 2 describes the
parameters for each accumulation measurement. While most of the data are measurements of accumulation between 1950
and present (~98%), to coincide with the time span of many regional climate models and reanalysis products, some ice core
records (~2%) are included dating back to 1800. The measurement methods include ice cores and/or boreholes, snow pits,
140 radar isochrons, and stake measurements. Arctic data are predominantly from ice cores and stake measurements and include
one radar transect in southeast Greenland (Bolzan and Strobel, 1999(a-g); Bolzan and Strobel, 2001(a,b); Mosley-Thompson
and others, 2001; Dibb and Fahnestock, 2004; Miège and others, 2013). The Antarctic data are predominantly from ice cores
and include two radar transects, one in West Antarctica and one in East Antarctica (Spikes and others, 2005; Banta and others,
2008; Ferris and others, 2011; Verfaillie and others, 2012; Burgener and others, 2013; Mayewski and others, 2013; Medley
145 and others, 2013). In most instances accumulation was provided in the original measurement, however, the Summit Station,
Greenland bamboo forest measurements consist of weekly surface height change at 100 stakes along with snow density (Dibb
and Fahnestock, 2004). We multiplied the height change by the coincident snow density and averaged across all stakes to get
accumulation measurements for SUMup. Similarly, the Bolzan and Strobel data (1999 a-g; 2001 a,b) provided a snow pit
depth, year, and density that were converted to accumulation. Most of the accumulation measurements are annually resolved
with the major exceptions being the radar data which is approximately decadal and bamboo forest data which is approximately
150 monthly.

Table 2

2.3.3 Snow Depth on Sea Ice

155 The snow depth on sea ice subdataset is the sparsest within SUMup with ~14,000 point measurements. Table 3 describes the
parameters for each snow depth measurement. The measurement methods include rulers and magnaprobos. The Arctic
measurements are from the Finnish Meteorological Institute Sea Ice Observers data from 1990-2012 (Eero Rinne, personal
communication). The Antarctic observations are from the Sea Ice Mass Balance in the Antarctic (SIMBA) dataset which was
conducted from the research vessel/ice breaker *N.B. Palmer* in September and October 2007 in the Bellingshausen Sea (Lewis
et al, 2011). Although this subdataset is rather small, we note that a large, standardized dataset of radar-derived snow depth
on sea ice is available through the IceBridge Sea Ice Freeboard, Snow Depth and Thickness product (Krutz and others, 2017)
160 as well as other products based off of the IceBridge Snow Radar (Kwok and Others, 2017). Because the IceBridge radar-
derived snow depths are too large and numerous they are no longer included in the SUMup dataset and were removed from
the July 2015 SUMup dataset for the current July 2017 SUMup dataset release.

Table 3

165 3. Data Analysis

The goal of the SUMup dataset is that it can be broadly used by the scientific community for a variety of research studies.
Tables 4 and 5 provide the basic descriptive statistics for each subdataset for the Arctic and Antarctic, respectively. These
tables provide a coarse overview of the data, however, when using the SUMup datasets subsetting by location, time, depth, etc
will likely be required for specific applications. The minimum value for accumulation in the Arctic is -0.004 m WE/a which
170 represents an ablation value from monthly Bamboo Forest measurements at Summit Station, Greenland. In total, there are 4
months with small negative accumulation (or ablation) measurements from Summit Station.

Field data collected over the vast polar regions have spatial and temporal sampling bias, as the time, cost and logistics to
systematically sample these regions is unreasonable. We describe the SUMup dataset here to elucidate possible bias. All the
175 measurements in SUMup, with the exception of one location, were collected during the the spring/summer season for that
polar region, roughly April through August for the Arctic and October through February for the Antarctic. Summit Station,



180 Greenland, the only GrIS station with year-round operations, is the an exception in the dataset where temporally-consistent, year-round measurements are taken. Below, we summarize the spatial and temporal distributions of the SUMup dataset by subdatasets. For the two largest subdatasets, snow density and accumulation we present preliminary analysis over the GrIS (Sect. 3.4). This analysis is meant to be an introduction to the dataset and is not exhaustive. We encourage the community to continue to use and more fully exploit this dataset.

Table 4

Table 5

185

3.1 Snow Density

Over 830,000 measurements were compiled of snow/firn density that cover ~280 sites in the Arctic and ~50 sites across Antarctica (Fig. 1). The majority of these measurements come from snow pits and ice cores on the GrIS and AIS, however, there are 7 locations of snow density measurements on sea ice in the Bellinghausen Sea. Here, we include only on the ice sheet measurements.

190

The Arctic snow density measurements, all on the GrIS, contain 94% of the measurements with <1% coming from Summit Station. The remaining 6% of the measurements come from the Antarctic. The density subdataset is dominated (97% of data) by high-depth resolution data (mm-scale for ~100 meters) from, X-ray microfocus computer tomography, neutron density methods and gamma-ray attenuation measurements taken on cores or in boreholes at 22 locations in Northern Greenland and on one core from West Antarctica. (Wilhelms, 2000a,b;c;d; Miller and Schwager, 2000a,b; Schaller and others, 2016; Kreutz et al, 2011). Because of the high-depth resolution of these data, compared to more typical density measurements at centimeter and meter scale, these data saturate histogram representations of this subdataset. For this reason we have removed these data from the following analysis to provide a more realistic overview of the fraction of the density measurements taken throughout time and space.

195

200

Figure 2

Figure 2 provides histograms showing the fraction of density measurements taken by year for Antarctica, Greenland excluding Summit Station and for Summit Station. (Summit Station was defined as a bounding box of 72N to 73N and 38W to 39W.) Summit Station measurements are plotted separately because this unique site provides the only location on the GrIS with year round measurements over multiple years. The histograms for the Antarctic and Greenland show sporadic spikes through time related to major collection campaigns. Antarctic density measurements peak in the early and late decade from 2000 to 2010 related to US ITASE traverses conducted in West and East, Antarctica, respectively (Mayewski and others, 2013). Greenland measurements peak in the early 1950's with measurements from Benson's traverses, are relatively stable through the 1990's related to the activities surrounding GISP2 and PARCA ice cores (Alley, 1999; Mosley-Thompson and others, 2001), and peak in the early 2010's with the Greenland Inland Traverse and the Arctic Circle Traverse cores (Miège and others, 2013; Hawley and others, 2014). Measurements from Summit station steadily increase in time from 1987 to 2014 with a slight peak in the late 1990's and early 2000's related to additional data provided by Dibb and Fahnestock (2004) and Alley (1999) in that year.

205

210

215

Figure 3

Figure 3 provides an overview of the distributions of depths sampled by the density subdataset. Overall, the number of measurements decrease with depth. The Antarctic data decreases more uniformly with depth which is related to the larger number of 50-100 m ice cores. The Greenland data decreases at a constant rate to 5 meters but has very few measurements below 25 m demonstrating the large number of shallow cores (~20 m) collected across Greenland. At Summit Station, the

220



225 majority of the measurements are taken above 1 meter as a result of systematic tasking to dig ~1 m snow pits at approximately
monthly intervals since 2003. The deep 100 m plus measurements at Summit come from the GISP2 ice core (Alley, 1999).

3.2 Accumulation

230 The ~230,000 measurements of accumulation rate over land ice were taken at ~30 locations in Antarctica, and ~35 locations
in Greenland. These include two radar traverses that span several hundreds of km in Antarctica, and a 75-km radar traverse in
Southeast Greenland. (Fig. 1).

235 62% of the accumulation measurements are from the Arctic, all within Greenland, with <1% of the overall measurements
coming from Summit Station. The Antarctic contains the remaining 38% of the measurements. The accumulation subdataset
is dominated (97% of data) by high-spatial resolution (10's of m) radar accumulation data taken from 3 ice sheet transects
(Fig. 1). These data saturate histogram representations of this subdataset and have been removed from the following analysis
to provide a more realistic overview of the fraction of accumulation measurements taken throughout time and space.

240 Figure 4 provides histograms showing the fraction of accumulation measurements taken by year for Antarctica, Greenland
excluding Summit Station, and for Summit Station. Year, in this case, is defined as the year in which the ice core, snow pit,
etc was collected/dug. The histograms for the Antarctic and Greenland show sporadic spikes through time related to major
collection campaigns, similar to, yet more exaggerated than, the density subdataset. Antarctic measurements peak in the early
2000's when US ITASE ice cores were collected in West Antarctica (Mayewski and others, 2013). Greenland accumulation
measurements peak in the late 1980's with ice cores preparing for the GISP2 core and in the late 1990's when the PARCA ice
cores (Mosley-Thompson and others, 2001) were collected. Summit Station has a constant monthly collection of accumulation
245 measurements from August 2000 to August 2002 from the monthly Bamboo Forest measurements (Dibb and Fahnestock,
2004) and represents the only year-round collection of accumulation in the SUMup dataset.

Figure 4

250 While understanding the date when accumulation measurements are taken is important it is arguably more important to
understand the years sampled by the ice core or snow pit. Figure 5 provides the distribution of years when annual accumulation
was measured from 1950 to present. Antarctica has a relatively even distribution of accumulation measurements until 2000
when the number of samples decreases. This decrease is due to the fact that many of the cores collected by US ITASE from
2006-2008 in East Antarctica could not be dated to determine accumulation and also shows that most of the ice cores collected
255 date back to 1950 or later. The Greenland accumulation measurements peak between 1980 and 2000. The mostly shallow ice
cores in Greenland, and relatively higher accumulation rates compared to Antarctica, cause there to be less data from 1950 to
1980 in the ice cores. The sharp decline in the 2000's is due to a lack of coring efforts that occurred during that decade in
Greenland. Summit Station has a consistent year round sampling of accumulation from the 2003 from the Bamboo Forest.
These systematic measurements significantly outnumber the single measurements per year collected from ice cores at Summit
260 Station that sample the decades previous to 2000.

Figure 5

3.3 Snow Depth on Sea Ice

265 The ~14,000 measurements of snow depth on sea ice are from 7 locations off of the Antarctic Peninsula in the Bellingshausen
sea (Fig. 1) and ~40 locations directly off the coast of Finland in the Baltic sea. The Arctic represents 78% of the measurements
and the Antarctica represents the remaining 22%. The Finnish Meteorological Institute Sea Ice Observers data (Eero Rinne,
personal communication) spans from from 1990-2012 (Fig. 6) while all the Antarctic data is all from 2007.

Figure 6



270 3.4 Preliminary Analysis over the Greenland Ice Sheet

3.4.1 Density distributions with Elevation, Latitude and Temperatures

275 Recent warming over the GrIS, including a melt event in 2012 that covered nearly the entire surface (Nghiem and others, 2012), has increased both snow density and snow accumulation in recent decades (e.g. Morris and Wingham, 2014; Machguth and others, 2016; Overly and others, 2016). Improved measurements, or models, of density and its evolution with time are needed to reduce uncertainties when converting altimetry measurements into total ice sheet mass balance using altimetry (e.g. Zwally and Li, 2002; Shepherd and others, 2012) and for converting radar isochrons into measurements of accumulation (e.g. Koenig and others, 2016). Many models use mean annual temperature and accumulation to model the spatial and temporal evolution of density (e.g. Harron and Langway, 1982; Reeh and others, 2005; Kuipers Munneke and others 2015). Some studies, however, show that density models generally underestimate surface (<1 m depth) density measurements (Koenig et al., 2016) while other studies point to the importance of the surface boundary condition for density models when comparing to measurements (Kuipers Munneke and others, 2015; Bellaire and others, 2017). Fausto and others (in revision) suggest two new snow surface (0-10 cm) density parameterizations, derived from a set of observations, using mean annual temperature and elevation to help modeling studies set surface boundary conditions. Here, we look more closely at the density and accumulation measurements within the SUMup dataset over the GrIS and their sampling distributions with respect to temperature, elevation and latitude.

Figure 7

290 Figure 7 shows the distribution of density measurements with elevation and latitude compared to the total distribution of elevations and latitudes for the entire GrIS derived from the Bamber 5 km Greenland digital elevation model (DEM; Bamber, 2001). Figure 7 uses similar graphing techniques to those of Fausto and others (in revision) to clearly show collection bias in the observation dataset. If there were no sampling bias, the fraction of measurements would be similar to the fraction of values from the DEM. This is not the case. For elevation (Fig. 7A) we see that elevations below 3000 m are undersampled, with the exception of the 1750-2000 m bin, and elevations above 3000 m are largely oversampled. The measurements are therefore biased to higher, inland elevations which, if averaged, would likely cause a low bias in sampled densities. Figure 7B shows that our dataset is sampled best over central Greenland. More measurements are required from lower elevations and southern (< 70N) and northern (> 78N) latitudes to fill the gaps in the current dataset and reduce spatial bias.

300

Figure 8

305 Because mean annual temperature is a parameter often used to model density (e.g. Harron and Langway, 1982; Reeh and others; 2005), Figure 8 shows the distribution of density measurements in Greenland by the mean annual temperature estimated by the Modèle Atmosphérique Régional (MAR) model version 3.5 (Fettweis and others, 2013). The density measurements at each location were matched to a National Centers for Environmental Prediction–National Center for Atmospheric Research Reanalysis version 1 (NCEP-NCARv1) forced MAR 3.5 simulation (run from 1948–2015) to find the mean annual 3 m air temperature for the year the measurement was taken. The NCEP-NCAR forcing was chosen because it is more reliable than ERA forcings (Fettweis et al, 2017). The red line in Fig. 8 shows the distribution of annual average temperatures (derived from 1990-2015) for the entire GrIS. Figure 8 clearly shows a preferential sampling of GrIS regions with lower temperatures. Cold temperatures (-20 deg C and below) are oversampled in the density dataset while temperatures above -14 deg C, which make up ~30% of the GrIS, make up less than 5% of the sampled densities. As with elevation, the density sampling distribution by mean annual temperatures likely results in a low density bias when trying to characterize the entire GrIS. In general, the density measurements in SUMup across the GrIS oversample cooler, inland regions and undersample coastal, warmer regions.

315

Figure 9



320 Figure 9 plots all sites in Greenland with density coincidentally sampled to depths of 10, 25, 50 and 100 cm compared to the mean annual temperature. No clear relationship (Pearson Correlation coefficient, $R^2 = 0$ to 0.057) between mean annual temperature and density is seen in our data until ~ 1 m depth ($R^2=0.255$) where higher temperatures correspond to higher density. This results suggests that in the top 1 m of snow/firn on the GrIS, in the colder, more inland areas, temperature may not be the primary process leading to densification. Solar radiation and wind processes (e.g. Liston and others, 2007) are likely important in these region and require snow density models that account for these processes. Due to the spatial sampling bias in this dataset, melt processes are likely not a primary processes in determining snow density for these measurements, however, melt processes will contribute more in the future (Nghiem and others, 2012; McGrath and others, 2015)

325

Figure 10

330 Figure 10 compares the observed snow densities to modeled densities from MAR 3.7 at 10, 20, 50 and 100 cm. In all cases the MAR model, on average, underestimates the near-surface snow density. The root mean-square error (RMSE) values are all between 0.07-0.08 indicating a small variance and generally good fit of the model. The mean biases are all negative values between -0.05 and -0.04 in agreement with the model underestimating the “true” value of the observations. Similar results were found by Koenig and others (2016) using a smaller subset of the SUMup data and holds with the newly added measurements. This suggests that snow on the ground is densifying at a faster rate in the colder, more inland locations of the GrIS, than the MAR model is predicting in the top meter.

335

3.4.2 Accumulation distributions with Elevation and Latitude

340 Snowfall over the GrIS can also be parameterized by elevation and latitude. Figures 11 show the distributions of the accumulation measurements over the GrIS by elevation and Latitude. As with the density measurements the accumulation measurements all come from high elevations on the GrIS (>1750 m) with the highest elevations (>300 m) being largely oversampled. The sampling across latitudes is the most evenly distributed, however, latitudes above 78 N represent a gap in the dataset. We do not compare the accumulation subdataset with mean annual temperatures here. Because each year of accumulation has a different mean annual temperature associated with it, we deem it beyond the scope of this analysis and suggest this as a future study that could be researched with the SUMup dataset.

345

Figure 11

3.4.3 Year round density and accumulation measurements from Summit Station.

350 Summit Station is the only site in the dataset, and on the GrIS, that has been systematically sampled for density and accumulation on a nearly monthly basis. Hence, it is the only location on the GrIS to watch the long-term, decadal, seasonal evolution of snow surface density. Figure 12 shows the seasonal trends of the mean surface density to depths of 10 cm, 25 cm 50 cm and 100 cm. A seasonal cycle is evident in the 10 cm and 25 cm depth mean densities with a decrease (trough) in density in late summer (August/September) and an increase (peak) in April. The decrease in summer density is likely due to surface hoar, a low density snow crystal, formation that is well known to form at Summit Station in the Summer when wind speeds are low and humidity relatively high (e.g. Alley and other, 1990; Albert and Schultz, 2002; Dibb and Fahnestock, 2004). As wind speed increase and water vapor decreases in the winter the surface snow increases in density. The seasonal signal in density is damped out by 1 m at Summit Station. Figure 12 also shows larger natural variability in average density measurements in the top 50 cm compared to the top 100 cm. This is expected as the deeper snow is more insulated from atmospheric and radiative processes in this dry-snow-zone location.

360

Figure 12



365 Figure 13 show the seasonal cycle of accumulation at Summit Station. Accumulation is highly variable with slightly lower values in the early Summer months (May/June/July). Dibb and Fahnestock (2004) also showed a similar trend in stake measurements and Summit Station from just 2 years of data and explained that the summer season may not actually be seeing a decrease in accumulation but that thinning layers, densification, may be causing the stake measurements to not rise as much in the summertime compared to the wintertime when a snowfall event occurs. Determining if there is a true decrease in summer accumulation or increase in snow/firn compaction rate at Summit Station requires additional research.

370 Figure 13

5. Data Availability

The SUMup dataset is currently available through the Arctic Data Center. It hosts our three subdatasets in both csv and netcdf formats along with metadata files to further explain the methods and citations. The dataset will be updated annually.

375 6. Discussion and Conclusion

380 We present and describe the SUMup dataset, an expandable, community-based dataset of field measurements of surface mass balance components that is consistent in format, properly described through metadata, and publically available. The subdatasets include compiled measurements of snow/firn density, accumulation on land ice, and snow depth on sea ice from the Arctic and Antarctica.

385 Though the SUMup dataset likely represents the largest of its kind, the measurements over the GrIS and AIS are sporadic in time and space, peaking during specific field campaigns and lapsing in between. This sampling strategy makes monitoring change with and understanding processes from field measurements difficult, especially for parameters like density and accumulation, that change with both seasonal and climatic atmospheric condition. Overall, there are gaps in density and accumulation data from ~2000 forward and from locations on the periphery of the ice sheets. While there currently is a temporal gap in the most recent decades we note that the GreenTRaCs traverses have collected cores across the GrIS in 2016 and 2017, including at previous PARCA sites. In the future, once processed, these cores will be help fill some of the time gap for the GrIS (R. Hawley, personal communication).

390 Density and accumulation measurements of the GrIS oversample cooler, inland regions and undersample coastal, warmer regions. Oversampling these regions may lead to an underestimation of the total average surface density, especially in the summer season, when the measurements are undersampling regions with significant melt processes that increase density. No clear relationship between mean annual temperature and density is seen in the data until a depth of 1 m where a relationship between higher temperatures and increased density is observed. This suggests that additional parameters, such as wind speed and radiative balance, should be considered when modeling density for the GrIS at SUMup density locations and depths above 50 cm. MAR estimated densities are lower than the SUMup measurements in the top 1 m. Possible causes for the underestimation include that the atmospheric snowfall is preferentially low, that the initial density of freshly fallen snow is underestimated, or that the surface snow on the ground densifies at a quicker rate than modelled by MAR, due to radiative or wind-driven processes in these mostly dry-snow zone locations. Summit Station, Greenland is the only location with year-round density and accumulation measurements in the dataset, and on the GrIS, and seasonal cycles are evident in accumulation rate and density for depths above 50 cm.

405 We encourage the cryospheric community to contribute additional field data to the SUMup dataset. We also encourage the cryospheric community, including modelers and remote sensors to use this dataset for model validation for surface mass



balance and satellite- or airborne-sensor algorithm development. SUMup is a dynamic, living dataset and is expected to be expanded and released annually.

Author Contributions:

410 LM compiled the SUMup dataset into the July 2017 dataset, developed the metadata and reformatted the dataset. She made all figures for this paper and co-wrote the paper. LK co-wrote this paper and developed the first SUMup dataset 2013. PA helped with the development of the SUMup dataset, performed the initial comparison of the SUMup data to the MAR model and contributed to the writing of this paper.

415 Competing Interests:

The authors declare that they have no conflict of interest.

Acknowledgements:

420 LM and LK acknowledge the National Science Foundation grant PLR 1603407 for funding this work. *Publication of this article was funded by the University of Colorado Boulder Libraries Open Access Fund.*

References:

- Albert M (2007) Snow and Firn Permeability: Characteristics of Snow Megadunes and their Potential Effects on Ice Core Interpretation. (doi:10.7265/N5639MPD)
- 425 Albert MR and Shultz EF (2002) Snow and firn properties and air-snow transport processes at Summit, Greenland. *Atmospheric Environment* **36**, 2789–2797 (doi:10.1016/S1352-2310(02)00119-X)
- Alley RB (1999) GISP2 Stratigraphy. (doi:10.1594/PANGAEA.56103)
- Alley RB, Saltzman ES, Cuffey KM and Fitzpatrick JJ (1990) Summertime formation of Depth Hoar in central Greenland. *Geophys. Res. Lett.* **17**(13), 2393–2396 (doi:10.1029/GL017i013p02393)
- 430 van Angelen JH, M. Lenaerts JT, van den Broeke MR, Fettweis X and van Meijgaard E (2013) Rapid loss of firn pore space accelerates 21st century Greenland mass loss: 21ST CENTURY GREENLAND SMB. *Geophysical Research Letters* **40**(10), 2109–2113 (doi:10.1002/grl.50490)
- Arthern RJ, Winebrenner DP and Vaughan DG (2006) Antarctic snow accumulation mapped using polarization of 4.3-cm wavelength microwave emission. *J. Geophys. Res.* **111**(D6), D06107 (doi:10.1029/2004JD005667)
- 435 Bales, R. C., McConnell, J. R., Mosley-Thompson, E., & Csatho, B. (2001). Accumulation over the Greenland ice sheet from historical and recent records. *Journal of Geophysical Research: Atmospheres*, *106*(D24), 33813-33825.
- Baker I (2016) NEEM Firn Core 2009S2 Density and Permeability. (doi:10.18739/A2Q88G)
- Bamber J. (2001) Greenland 5 km DEM, Ice Thickness, and Bedrock Elevation Grids. Boulder, Colorado USA: NASA DAAC at the National Snow and Ice Data Center.
- 440 Banta JR, McConnell JR, Frey MM, Bales RC and Taylor K (2008) Spatial and temporal variability in snow accumulation at the West Antarctic Ice Sheet Divide over recent centuries. *Journal of Geophysical Research* **113**(D23) (doi:10.1029/2008JD010235)
- Bellaire S, Proksch M, Schneebeli M, Niwano M and Steffen K (2017) Measured and Modeled Snow Cover Properties across the Greenland Ice Sheet. *The Cryosphere Discuss.* **2017**, 1–22 (doi:10.5194/tc-2017-55)
- 445 Benson C (2013) Greenland Snow Pit and Core Stratigraphy (Analog and Digital Formats) Boulder, Colorado USA: National Snow and Ice Data Center.
- Benson C (2017) Greenland Snow it and Core Stratigraphy. Carl S. Benson Collection. Coll. 2010011. Roger G. Barry Archives and Resource Center. National Snow Data Center. [Accessed June 2017].
- Bolzan JF and Strobel M (1999a) Oxygen isotope data from snowpit at GISP2 Site 13. (doi:10.1594/PANGAEA.55510)
- 450 Bolzan JF and Strobel M (1999b) Oxygen isotope data from snowpit at GISP2 Site 15. (doi:10.1594/PANGAEA.55511)
- Bolzan JF and Strobel M (1999c) Oxygen isotope data from snowpit at GISP2 Site 31. (doi:10.1594/PANGAEA.55512)
- Bolzan JF and Strobel M (1999d) Oxygen isotope data from snowpit at GISP2 Site 37. (doi:10.1594/PANGAEA.55513)
- Bolzan JF and Strobel M (1999e) Oxygen isotope data from snowpit at GISP2 Site 51. (doi:10.1594/PANGAEA.55514)
- Bolzan JF and Strobel M (1999f) Oxygen isotope data from snowpit at GISP2 Site 57. (doi:10.1594/PANGAEA.55515)
- 455 Bolzan JF and Strobel M (1999g) Oxygen isotope data from snowpit at GISP2 Site 73. (doi:10.1594/PANGAEA.55516)



- Bolzan JF and Strobel M (2001a) Oxygen isotope data from snowpit at GISP2 Site 44. (doi:10.1594/PANGAEA.59995)
- Bolzan JF and Strobel M (2001b) Oxygen isotope data from snowpit at GISP2 Site 571. (doi:10.1594/PANGAEA.59996)
- Broeke M van den, Bamber J, Ettema J, Rignot E, Schrama E, Berg WJ van de, Meijgaard E van, Velicogna I and Wouters B (2009) Partitioning Recent Greenland Mass Loss. *Science* **326**(5955), 984–986 (doi:10.1126/science.1178176)
- 460 Burgener L, Rupper S, Koenig L, Forster R, Christensen WF, Williams J, Koutnik M, Miège C, Steig EJ, Tingey D, Keeler D and Riley L (2013) An observed negative trend in West Antarctic accumulation rates from 1975 to 2010: Evidence from new observed and simulated records: West Antarctic Accumulation Rate Trends. *Journal of Geophysical Research: Atmospheres* **118**(10), 4205–4216 (doi:10.1002/jgrd.50362)
- 465 Burgess EW, Forster RR, Box JE, Mosley-Thompson E, Bromwich DH, Bales RC and Smith LC (2010) A spatially calibrated model of annual accumulation rate on the Greenland Ice Sheet (1958–2007). *J. Geophys. Res.* **115**(F2), F02004 (doi:10.1029/2009JF001293)
- Chellman (2016) Core Atmospheric and Snow Measurements at Summit Greenland Environmental Observatory: Snow Pit. (doi:10.18739/A2888F)
- Conway H (2003) Roosevelt Island Ice Core Density and Beta Count Data. (doi:10.7265/N55718ZW)
- 470 Dibb JE and Fahnestock M (2004) Snow accumulation, surface height change, and firn densification at Summit, Greenland: Insights from 2 years of in situ observation. *J. Geophys. Res.* **109**(D24), D24113 (doi:10.1029/2003JD004300)
- Dibb JE, Whitlow SI and Arsenaault M (2007) Seasonal variations in the soluble ion content of snow at Summit, Greenland: Constraints from three years of daily surface snow samples. *Atmospheric Environment* **41**(24), 5007–5019 (doi:10.1016/j.atmosenv.2006.12.010)
- 475 Dibb JE (2017) Personal Communication
- Enderlin EM, Howat IM, Jeong S, Noh M-J, van Angelen JH and van den Broeke MR (2014) An improved mass budget for the Greenland ice sheet. *Geophys. Res. Lett.* **41**(3), 2013GL059010 (doi:10.1002/2013GL059010)
- Ferris DG, Cole-Dai J, Reyes AR and Budner DM (2011) South Pole ice core record of explosive volcanic eruptions in the first and second millennia A.D. and evidence of a large eruption in the tropics around 535 A.D. *Journal of Geophysical Research* **116**(D17) (doi:10.1029/2011JD015916)
- 480 Fettweis X, Box JE, Agosta C, Amory C, Kittel C, Lang C, van As D, Machguth H and Gallée H (2017) Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model. *The Cryosphere* **11**(2), 1015–1033 (doi:10.5194/tc-11-1015-2017)
- 485 Hall DK, Comiso JC, DiGirolamo NE, Shuman CA, Box JE and Koenig LS (2013) Variability in the surface temperature and melt extent of the Greenland ice sheet from MODIS. *Geophys. Res. Lett.* **40**(10), 2114–2120 (doi:10.1002/grl.50240)
- Hastings, M. and Dibb, J. (2017) Personal Communication
- Hawley RL, Courville ZR, Kehrl LM, Lutz ER, Osterberg EC, Overly TB and Wong GJ (2014) Recent accumulation variability in northwest Greenland from ground-penetrating radar and shallow cores along the Greenland Inland Traverse. *Journal of Glaciology* **60**(220), 375–382 (doi:10.3189/2014JoG13J141)
- 490 Herron SL and Langway CC (1982) A Comparison of Ice Fabrics and Textures at Camp Century, Greenland and Byrd Station, Antarctica. *Annals of Glaciology* **3**, 118–124 (doi:10.1017/S0260305500002639)
- Koenig L, Box J and Kurtz N (2013) Improving Surface Mass Balance Over Ice Sheets and Snow Depth on Sea Ice. *Eos Trans. AGU* **94**(10), 100–100 (doi:10.1002/2013EO100006)
- Koenig L and Brucker L (2011) Satellite-Era Accumulation Traverse 2011 (SEAT11) snowpit density data.
- 495 Koenig LS, Ivanoff A, Alexander PM, MacGregor JA, Fettweis X, Panzer B, Paden JD, Forster RR, Das I, McConnell JR, Tedesco M, Leuschen C and Gogineni P (2016) Annual Greenland accumulation rates (2009–2012) from airborne snow radar. *The Cryosphere* **10**(4), 1739–1752 (doi:10.5194/tc-10-1739-2016)
- Koenig LS, Miège C, Forster RR and Brucker L (2014) Initial in situ measurements of perennial meltwater storage in the Greenland firn aquifer: Measurements of Greenland Aquifer. *Geophysical Research Letters* **41**(1), 81–85 (doi:10.1002/2013GL058083)
- 500 Kreutz K (2011) Microparticle, Conductivity, and Density Measurements from the WAIS Divide Deep Ice Core, Antarctica. (doi:10.7265/N5K07264)
- Kuipers Munneke P, Ligtenberg SRM, Noël BPY, Howat IM, Box JE, Mosley-Thompson E, McConnell JR, Steffen K, Harper JT, Das SB and van den Broeke MR (2015) Elevation change of the Greenland Ice Sheet due to surface mass balance and firn processes, 1960–2014. *The Cryosphere* **9**(6), 2009–2025 (doi:10.5194/tc-9-2009-2015)
- 505




- Kurtz N (2012) IceBridge Sea Ice Freeboard, Snow Depth, and Thickness, Version 1. (doi:10.5067/7XJ9HRV50O57)
- 510 Kwok R, Kurtz NT, Brucker L, Ivanoff A, Newman T, Farrell SL, King J, Howell S, Webster MA, Paden J, Leuschen C, Macgregor JA, Richter-Menge J, Harbeck J and Tschudi M (2017) Inter-comparison of snow depth retrievals over Arctic sea ice from radar data acquired by Operation IceBridge. *The Cryosphere Discuss.* **2017**, 1–37 (doi:10.5194/tc-2017-103)
- Lewis MJ, Tison JL, Weissling B, Delille B, Ackley SF, Brabant F and Xie H (2011) Sea ice and snow cover characteristics during the winter–spring transition in the Bellingshausen Sea: An overview of SIMBA 2007. *Deep Sea Research Part II: Topical Studies in Oceanography* **58**(9–10), 1019–1038 (doi:10.1016/j.dsr2.2010.10.027)
- 515 Liston GE, Haehnel RB, Sturm M, Hiemstra CA, Berezovskaya S and Tabler RD (2007) Instruments and Methods Simulating complex snow distributions in windy environments using SnowTran-3D. *Journal of Glaciology* **53**(181), 241–256 (doi:10.3189/172756507782202865)
- 520 Machguth H, Thomsen HH, Weidick A, Ahlstrøm AP, Abermann J, Andersen ML, Andersen SB, Bjørk AA, Box JE, Braithwaite RJ, Bøggild CE, Citterio M, Clement P, Colgan W, Fausto RS, Gleie K, Gubler S, Hasholt B, Hynek B, Knudsen NT, Larsen SH, Mernild SH, Oerlemans J, Oerter H, Olesen OB, Smeets CJPP, Steffen K, Stober M, Sugiyama S, As DV, Broeke MRVD and Wal RSWVD (2016) Greenland surface mass-balance observations from the ice-sheet ablation area and local glaciers. *Journal of Glaciology* **62**(235), 861–887 (doi:10.1017/jog.2016.75)
- Mayewski P and Whitlow S (2009a) Regional Survey of Greenland, 1988 - Snow Pit Data, Version 1.0. (doi:10.5065/D6154F6J)
- 525 Mayewski P and Whitlow S (2009b) Snow Pit and Ice Core Data from Southern Greenland, 1984, Version 1.0. (doi:10.5065/D6S180MH)
- Mayewski P and Whitlow S (2009c) Snow Pit Data from Greenland Summit, 1987, Version 1.0. (doi:10.5065/D63X84RQ)
- Mayewski P and Whitlow S (2009d) Snow Pit Data from Greenland Summit, 1989 to 1993, Version 1.0. (doi:10.5065/D6NP22KX)
- 530 McGrath D, Colgan W, Bayou N, Muto A and Steffen K (2013) Recent warming at Summit, Greenland: Global context and implications. *Geophys. Res. Lett.* **40**(10), 2091–2096 (doi:10.1002/grl.50456)
- Medley B, Joughin I, Das SB, Steig EJ, Conway H, Gogineni S, Criscitiello AS, McConnell JR, Smith BE, van den Broeke MR, Lenaerts JTM, Bromwich DH and Nicolas JP (2013) Airborne-radar and ice-core observations of annual snow accumulation over Thwaites Glacier, West Antarctica confirm the spatiotemporal variability of global and regional atmospheric models. *Geophys. Res. Lett.* **40**(14), 3649–3654 (doi:10.1002/grl.50706)
- 535 Miège C, Forster RR, Box JE, Burgess EW, McConnell JR, Pasteris DR and Spikes VB (2013) Southeast Greenland high accumulation rates derived from firn cores and ground-penetrating radar. *Annals of Glaciology* **54**(63), 322–332 (doi:10.3189/2013AoG63A358)
- Miller H and Schwager M (2000a) Density of ice core ngt37C95.2 from the North Greenland Traverse. (doi:10.1594/PANGAEA.57798)
- 540 Miller H and Schwager M (2000b) Density of ice core ngt42C95.2 from the North Greenland Traverse. (doi:10.1594/PANGAEA.57655)
- Mock SJ (1967a) Calculated Patterns of Accumulation on the Greenland Ice Sheet. *Journal of Glaciology* **6**(48), 795–803 (doi:10.1017/S0022143000020104)
- 545 Mock SJ (1967b) Accumulation Patterns On The Greenland Ice Sheet. CRREL-RR-233. Cold Regions Research And Engineering Lab Hanover NH
- Morris EM and Wingham DJ (2014) Densification of polar snow: Measurements, modeling, and implications for altimetry. *J. Geophys. Res. Earth Surf.* **119**(2), 349–365 (doi:10.1002/2013JF002898)
- 550 Mosley-Thompson E, McConnell JR, Bales RC, Li Z, Lin P-N, Steffen K, Thompson LG, Edwards R and Bathke D (2001) Local to regional-scale variability of annual net accumulation on the Greenland ice sheet from PARCA cores. *J. Geophys. Res.* **106**(D24), 33839–33851 (doi:10.1029/2001JD900067)
- Nghiem SV, Hall DK, Mote TL, Tedesco M, Albert MR, Keegan K, Shuman CA, DiGirolamo NE and Neumann G (2012) The extreme melt across the Greenland ice sheet in 2012. *Geophys. Res. Lett.* **39**(20), L20502 (doi:10.1029/2012GL053611)
- 555 Ohmura A (1991) ETH Greenland Expedition I, progress report no. 1, April 1989 to February, 1991. Tech. Report, Department of Geography, ETH Zürich, 108 pp.



- Ohmura A (1992) ETH Greenland Expedition II, progress Rep. 2, April 1991 to October, 1992. Tech. Report, Department of Geography, ETH Zürich, 94 pp.
- Ohmura A and Reeh N (1991) New precipitation and accumulation maps for Greenland. *Journal of Glaciology* **37**(125), 140–148 (doi:10.1017/S0022143000042891)
- 560 Overly TB, Hawley RL, Helm V, Morris EM and Chaudhary RN (2016) Greenland annual accumulation along the EGIG line, 1959–2004, from ASIRAS airborne radar and neutron-probe density measurements. *The Cryosphere* **10**(4), 1679–1694 (doi:10.5194/tc-10-1679-2016)
- Reeh N, Fisher DA, Koerner RM and Clausen HB (2005) An empirical firn-densification model comprising ice lenses. *Annals of Glaciology* **42**(1), 101–106 (doi:10.3189/172756405781812871)
- 565 Richter-Menge J, Overland JE and Mathis JT (2016) Arctic Report Card: Update for 2016 – LSU Law Center: Climate Change Law and Policy Project. <https://sites.law.lsu.edu/coast/2016/12/arctic-report-card-update-for-2016/>
- Schaller CF, Freitag J, Kipfstuhl S, Laepple T, Steen-Larsen HC and Eisen O (2016) A representative density profile of the North Greenland snowpack. *The Cryosphere* **10**(5), 1991–2002 (doi:10.5194/tc-10-1991-2016)
- Serreze MC and Francis JA (2006) The Arctic on the fast track of change. *Weather* **61**(3), 65–69 (doi:10.1256/wea.197.05)
- 570 Shepherd A, Ivins ER, A G, Barletta VR, Bentley MJ, Bettadpur S, Briggs KH, Bromwich DH, Forsberg R, Galin N, Horwath M, Jacobs S, Joughin I, King MA, Lenaerts JTM, Li J, Ligtenberg SRM, Luckman A, Luthcke SB, McMillan M, Meister R, Milne G, Mouginot J, Muir A, Nicolas JP, Paden J, Payne AJ, Pritchard H, Rignot E, Rott H, Sørensen LS, Scambos TA, Scheuchl B, Schrama EJO, Smith B, Sundal AV, Angelen JH van, Berg WJ van de, Broeke MR van den, Vaughan DG, Velicogna I, Wahr J, Whitehouse PL, Wingham DJ, Yi D, Young D and Zwally HJ (2012) A Reconciled Estimate of Ice-Sheet Mass Balance. *Science* **338**(6111), 1183–1189 (doi:10.1126/science.1228102)
- 575 US International Trans-Antarctic Scientific Expedition (US ITASE) Glaciochemical Data, Version 2- Mayewski, P. A. and D. A. Dixon. 2013. US International Trans-Antarctic Scientific Expedition (US ITASE) Glaciochemical Data. Version 2. [US_ITASE_Core Info-SWE-Density_2013.xlsx]. Boulder, Colorado USA: National Snow and Ice Data Center.
- V. Spikes GH (2005) US International Trans-Antarctic Scientific Expedition (US ITASE): GPR Profiles and Accumulation Mapping. (doi:10.7265/N5GH9FV6)
- 580 Vaughan DG, Bamber JL, Giovinetto M, Russell J and Cooper APR (1999) Reassessment of Net Surface Mass Balance in Antarctica. *J. Climate* **12**(4), 933–946 (doi:10.1175/1520-0442(1999)012<0933:RONSMB>2.0.CO;2)
- Vaughan DG, Marshall GJ, Connolley WM, Parkinson C, Mulvaney R, Hodgson DA, King JC, Pudsey CJ and Turner J (2003) Recent Rapid Regional Climate Warming on the Antarctic Peninsula. *Climatic Change* **60**(3), 243–274 (doi:10.1023/A:1026021217991)
- 585 Velicogna I, Sutterley TC and van den Broeke MR (2014) Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data. *Geophys. Res. Lett.* **41**(22), 8130–8137 (doi:10.1002/2014GL061052)
- Verfaillie D, Fily M, Le Meur E, Magand O, Jourdain B, Arnaud L and Favier V (2012) Snow accumulation variability derived from radar and firn core data along a 600 km transect in Adelie Land, East Antarctic plateau. *The Cryosphere* **6**(6), 1345–1358 (doi:10.5194/tc-6-1345-2012)
- 590 Wilhelms F (2000a) Density of ice core ngt03C93.2 from the North Greenland Traverse. (doi:10.1594/PANGAEA.56560)
- Wilhelms F (2000b) Density of ice core ngt06C93.2 from the North Greenland Traverse. (doi:10.1594/PANGAEA.57153)
- Wilhelms F (2000c) Density of ice core ngt14C93.2 from the North Greenland Traverse. (doi:10.1594/PANGAEA.56615)
- 595 Wilhelms F (2000d) Density of ice core ngt27C94.2 from the North Greenland Traverse. (doi:10.1594/PANGAEA.57296)
- Zwally HJ and Jun L (2002) Seasonal and interannual variations of firn densification and ice-sheet surface elevation at Greenland summit. *Journal of Glaciology* **48**, 199–207 (doi:10.3189/172756502781831403)

600

605



610

Figure 1 a) Measurement locations for accumulation (red), snow density (blue) or both (purple) for the SUMup dataset in the Arctic. Snow depth on sea ice data for the Arctic are not shown but include ~40 measurements taken off the coast of Finland in the Baltic Sea. b) Same as above but for the Antarctic. All locations not on the Antarctic ice sheet, those in the Bellingshausen Sea, include both snow density and snow depth on sea ice measurements.

620

625

630



635

Table 1: The parameters for each snow density measurement in the SUMup dataset with a brief description and the unit of measurement.

<u>Column</u>	<u>Description</u>	<u>Unit</u>
Date Taken	Date the data was taken	yyyymmdd
Latitude	Latitude of measurement	Decimal degree
Longitude	Longitude of measurement	Decimal degree
Start Depth	Top depth of the measurement in m from the snow/air interface (snow surface).	m
Stop Depth	Bottom depth of the measurement in m from the snow/air interface (snow surface).	m
Midpoint Depth	Midpoint depth of the measurement in m from the snow/air interface (snow surface).	m
Density	Snow density measurement	g/cm ³
Error	Uncertainty in density measurement	g/cm ³
Elevation	Elevation above sea level	m
Method	How the measurement was collected (see metadata for more details)	-
Citation	Cited source of data (see metadata for more details)	-

640

645

650

655



660

Table 2: The parameters for each snow accumulation measurement on land ice in the SUMup dataset with a brief description and the unit of measurement.

<u>Column</u>	<u>Description</u>	<u>Unit</u>
Date Taken	Date the data was taken	yyyymmdd
Latitude	Latitude of measurement	Decimal degree
Longitude	Longitude of measurement	Decimal degree
Start Year	First year of measurement if accumulation is not annual	year
End Year	Last year of measurement if accumulation is not annual	year
Year	Year of accumulation if accumulation is annual	year
Accumulation	Accumulation in m of water equivalent	m WE/a
Error	Uncertainty in measurement	m WE/a
Elevation	Elevation above sea level	m
Radar Horizontal Resolution	Horizontal resolution of radar data along track	m
Method	How the measurement was collected (see metadata for more details)	-
Name	Name of field campaign (see metadata for more details)	-
Citation	Cited source of data (see metadata for more details)	-

665

670

675

680



685 Table 3: The parameters for each snow depth on sea ice measurement in the SUMup dataset with a brief description and the unit of measurement

<u>Column</u>	<u>Description</u>	<u>Unit</u>
Date Taken	Date the data was taken	yyyymmdd
Latitude	Latitude of measurement	Decimal degree
Longitude	Longitude of measurement	Decimal degree
Distance Along Transect	Distance along a transect of in-situ snow depth measurements over sea ice from the initial Lat, Long. Used for snow-depth measurements where point by point Lat, Long was not recorded.	m
Snow Depth	Snow depth measurement	m
Snow Depth Error	Uncertainty in snow depth measurement	m
Density Taken	If density measurement was taken = 1, if no measurement =0.	-
Sea Ice Thickness	Sea ice thickness measurement	m
Sea Ice Thickness Error	Uncertainty in sea ice thickness measurement	m
Sea Ice Type	1=first year ice, 2=multilayer ice, -9999 = unknown	-
Sea Ice Freeboard	Sea Ice freeboard measurement	m
Sea Ice Freeboard Error	Uncertainty in sea ice freeboard	m
Snow Ice Thickness	Snow ice thickness measurement	m
Snow Ice Thickness Error	Uncertainty in snow ice thickness measurement	m
Radar Horizontal Resolution	Horizontal resolution of radar data along track	m
Method	How the measurement was collected (see metadata for more details)	-
Citation	Cited source of data (see metadata for more details)	-

690

695



Table 4: The descriptive statistics for the all of the Arctic measurements in SUMup including the minimum (min), maximum (max), mean, median, standard deviation (stdev) and number of measurements (N).

700

Statistic	Snow Density (g cm^{-3})	Accumulation on Land Ice (m WE/a)	Snow Depth on Sea Ice (m)
min	0.054	-0.004	0.000
max	0.970	2.257	0.500
mean	0.595	1.040	0.043
median	0.629	1.034	0.000
stdev	0.236	0.171	0.072
N	779439	144060	10979

705

710

715

720

725

730



735 Table 5: The descriptive statistics for the all of the Antarctic measurements in SUMup including the minimum (min), maximum (max), mean, median, standard deviation (stdev) and number of measurements (N).

Statistic	Snow Density(g cm ⁻³)	Accumulation on Land Ice (m WE/a)	Snow Depth on Sea Ice (m)
min	0.208	0.019	0.000
max	0.938	0.790	1.900
mean	0.789	0.129	0.479
median	0.841	0.132	0.430
stdev	0.130	0.055	0.363
N	51664	86509	3176

740


745

750

755

760

765



770

Figure 2. Histograms showing the date taken and associated fraction of the density dataset for A) Antarctica, B) Greenland excluding Summit Station, and C) Summit Station.

775

780

785

790

795




Figure 3. Histograms showing the fraction of the density dataset by mid-point sampling depths for A) Antarctica, B) Greenland excluding Summit Station and C) Summit Station.


800

805

810

815

820



825 Figure 4. Histograms showing the date taken and associated fraction of the accumulation dataset for each area examined: A) Antarctica, B) Greenland excluding Summit Station, and C) Summit Station.


830

835

840

845

850



855 Figure 5: Histograms showing the fraction of accumulation measurements by year for A) Antarctica, B) Greenland excluding
Summit Station and for C) Summit Station. Dates are only shown from 1950 forward.

860

865

870

875

880

885



890




Figure 6: Histogram showing the fraction of snow depth on sea ice measurements by year.

895


900

905

910



915



920 Figure 7: A) Histogram showing the fraction of density subset by elevation. Red line is the fraction of elevations for the
entire GrIS from Bamber DEM. B) Histogram showing the fraction of density dataset by latitude. Red line is the fraction of
925 latitudes for the entire GrIS using Bamber DEM.

925

930

935



940




Figure 8: Histogram showing the fraction of the density subdataset by modeled 3m annual air temperature. Red line shows 1990-2015 annual average MAR3.5 model 3m air temperature distribution for each grid cell across the ice sheet.

945


950

955

960



965



970 Figure 9. Scatterplot showing the MAR 3.5 modeled mean annual 3 m air temperature in the year the density was measured
971 compared to the mean density in top A) 10 cm, B) 25 cm, C) 50 cm, and D) 100 cm.

975

980



985

990




Figure 10: Scatterplot showing observed density (Y-axis) vs. MAR3.7 (x-axis) modelled density. One to one line in red. A) 10 cm B) 20 cm C) 50 cm D) 100 cm

995

1000



1005




Figure 11: A) Histogram showing the fraction of accumulation subset by elevation. Red line is the fraction of elevations for the entire GrIS from Bamber DEM. B) Histogram showing the fraction of accumulation dataset by latitude. Red line is the fraction of latitudes for the entire GrIS using Bamber DEM.

1015


1020

1025

1030



1035



1040


Figure 12: Plot of mean density (circle) and +/- 1 standard deviation (whiskers) for each month at Summit Station, Greenland for A) 10 cm b) 25 cm c) 50 cm d) 100 cm

1045

1050



1055



1060 Figure 13: Plot of mean accumulation (circle) and +/- 1 standard deviation (whiskers) for each month at Summit Station, Greenland.