

# Modern and Ancient Continental Shelf Anoxia

EDITED BY

**R. V. TYSON**

University of Newcastle upon Tyne  
UK

**T. H. PEARSON**

Scottish Environmental Advisory Services Ltd  
UK

1991

Published by

The Geological Society

London

# Contents

|                                                                                                                                                                                                                         |     |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| TYSON, R. V. & PEARSON, T. H. Modern and ancient continental shelf anoxia: an overview                                                                                                                                  | 1   |
| <b>Modern shelf anoxia</b>                                                                                                                                                                                              |     |
| BOESCH, D. F. & RABALAIS, N. N. Effects of hypoxia on continental shelf benthos: comparisons between the New York Bight and the Northern Gulf of Mexico                                                                 | 27  |
| RABALAIS, N. N., TURNER, R. E., WISEMAN, W. I. Jr. & BOESCH, D. F. A brief summary of hypoxia on the northern Gulf of Mexico continental shelf: 1985–1988                                                               | 35  |
| HARPER, D. E. Jr., MCKINNEY, L.D., NANCE, J. M. & SALZER, R. R. Recovery responses of two benthic assemblages following an acute hypoxic event on the Texas continental shelf, northwestern Gulf of Mexico              | 49  |
| VAN DER ZWAAN, G. J. & JORISSEN, F. J. Biofacial patterns in river-induced shelf anoxia                                                                                                                                 | 65  |
| MALONE, T. C. River flow, phytoplankton production and oxygen depletion in Chesapeake Bay                                                                                                                               | 83  |
| JUSTIĆ, D. Hypoxic conditions in the northern Adriatic Sea: historical development and ecological significance                                                                                                          | 95  |
| FAGANELI, J., PEZDIČ, J., OGORELEC, B., HERNDL, G. J. & DOLENEC, T. The role of sedimentary biogeochemistry in the formation of hypoxia in shallow coastal waters (Gulf of Trieste, northern Adriatic)                  | 107 |
| STACHOWITSCH, M. Anoxia in the Northern Adriatic Sea: rapid death, slow recovery                                                                                                                                        | 119 |
| ARNTZ, W. E., TARAZONA, J., GALLARDO, V. A., FLORES, L. A. & SALZWEDEL, H. Benthos communities in oxygen deficient shelf and upper slope areas of the Peruvian and Chilean Pacific coast, and changes caused by El Niño | 131 |
| EMEIS, K.-C., WHELAN, J. K. & TARAF, M. Sedimentary and geochemical expressions of oxic and anoxic conditions on the Peru shelf                                                                                         | 155 |
| BAILEY, G. W. Organic carbon flux and development of oxygen deficiency on the modern Benguela continental shelf south of 22°S: spatial and temporal variability                                                         | 171 |
| <b>Ancient shelf anoxia</b>                                                                                                                                                                                             |     |
| RHOADS, D. C., MULSOW, S. G., GUTSCHICK, R., BALDWIN, C. T., & STOLZ, J. F. The dysaerobic zone revisited: a magnetic facies?                                                                                           | 187 |
| SAVRDA, C. E. & BOTTJER, D. J. Oxygen-related biofacies in marine strata: an overview and update                                                                                                                        | 201 |
| CUOMO, M. C. & BARTHOLOMEW, P. R. Pelletal black shale fabrics: their origin and significance                                                                                                                           | 221 |
| BAIRD, G. C. & BRETT, C. E. Submarine erosion on the anoxic sea floor: stratigraphic, palaeoenvironmental and temporal significance of reworked pyrite-bone deposits                                                    | 233 |
| HECKEL, P. H. Thin widespread Pennsylvanian black shales of Midcontinent North America: a record of a cyclic succession of widespread pycnoclines in a fluctuating epeiric sea                                          | 259 |
| PIASECKI, S. & STEMMERIK, L. Late Permian anoxia in central East Greenland                                                                                                                                              | 275 |
| WIGNALL, P. B. & HALLAM, A. Biofacies, stratigraphic distribution and depositional models of British onshore Jurassic black shales                                                                                      | 291 |
| LITKE, R., BAKER, D. R., LEYTHAEUSER, D. & RULLKÖTTER, J. Keys to the depositional history of the Posidonia Shale (Toarcian) in the Hils Syncline, northern Germany                                                     | 311 |
| PRAUSS, M., LIGOUIS, B. & LUTERBACHER, H. Organic matter and palynomorphs in the 'Posidonienschiefer' (Toarcian, Lower Jurassic) of southern Germany                                                                    | 335 |

|                                                                                                                                                                                                            |     |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| BRUMSACK, H.-J. Inorganic geochemistry of the German 'Posidonia Shale': palaeoenvironmental consequences                                                                                                   | 353 |
| HUDSON, J. D. & MARTILL, D. M. The Lower Oxford Clay: production and preservation of organic matter in the Callovian (Jurassic) of central England                                                         | 363 |
| OSCHMANN, W. Distribution, dynamics and palaeoecology of Kimmeridgian (Upper Jurassic) shelf anoxia in western Europe                                                                                      | 381 |
| DOYLE, P. & WHITHAM, A. G. Palaeoenvironments of the Nordenskjöld Formation: an Antarctic Late Jurassic–Early Cretaceous black shale-tuff sequence                                                         | 397 |
| BRÉHÉRET, J.-G. Glauconitization episodes in marginal settings as echoes of mid-Cretaceous anoxic events in the Vocontian Basin (SE France)                                                                | 415 |
| KOUTSOUKOS, E. A. M., MELLO, M. R. & DE AZAMBUJA FILHO, N. C. Micropalaeontological and geochemical evidence of mid-Cretaceous anoxic dysoxic–palaeoenvironments in the Sergipe Basin, northeastern Brazil | 427 |
| VETŐ, I. & HETÉNYI, M. Fate of organic carbon and reduced sulphur in dysoxic–anoxic Oligocene facies of the Central Parathethys (Carpathian Mountains and Hungary)                                         | 449 |
| Index                                                                                                                                                                                                      | 461 |